

IV Antibiotic Administration at Home

1. Scrub injection cap.

- 15 seconds to scrub.
- 15 seconds to dry for alcohol wipe.

OR

- 5 seconds to scrub.
- 5 seconds to dry for Prevantix wipe.



2. Flush line with saline. Hold cap firmly when detaching syringe.



3. Scrub injection cap).



OR



- 15 seconds to scrub.
- 15 seconds to dry for alcohol wipe.

- 5 seconds to scrub.
- 5 seconds to dry for Prevantix wipe.

4. Attach antibiotic and unclamp tubing. Let infuse.



5. Scrub injection cap.



OR



- 15 seconds to scrub.
- 15 seconds to dry for alcohol wipe.

- 5 seconds to scrub.
- 5 seconds to dry for Prevantix wipe.

6. Flush line with saline (see step 2).



continue →

7. Scrub injection cap.



OR



- 15 seconds to scrub.
- 15 seconds to dry for alcohol wipe.
- 5 seconds to scrub.
- 5 seconds to dry for Prevantix wipe.

8. Flush with Heparin as directed.



9. Attach swab cap.



Please use the camera on your smartphone to scan the QR code and view related videos on our YouTube channel.



 **Children's Hospital
of The King's Daughters**

CHKD.org